

Claims Listng

1. (Currently Amended) A dosage form of a pharmaceutical composition which comprises an effective amount of a centrally-acting acetylcholinesterase inhibitor useful in the treatment of Alzheimer's disease selected from the group consisting of galanthamine, lycoramine, analogs of galanthamine and lycoramine rivastigmine said compounds having a half life of from one to eleven hours wherein ~~the~~ acetylcholinesterase inhibitor is formulated so as to delay its activity for a predetermined period of from four to twelve hours.
2. Canceled
3. (Previously presented) A dosage form of a pharmaceutical composition as claimed in claim 1 wherein the composition is formulated to delay the activity of the acetyl cholinesterase inhibitor for a period of from six to nine hours.
4. (Previously presented) A dosage form of a pharmaceutical composition as claimed in claim 1 wherein the composition is formulated to delay the activity of the acetyl cholinesterase inhibitor for a period of from eight to twelve hours.
5. (Original) A dosage form of a pharmaceutical composition as claimed in claim 1 wherein said acetylcholinesterase inhibitor has a duration of action of from 2 to 12 hours.
6. Canceled
7. (Original) A dosage form of a pharmaceutical composition as claimed in claim 1 wherein said acetylcholinesterase inhibitor is selected from the group consisting of galanthamine, lycoramine and analogs of said compounds wherein the methoxy group thereof is replaced by another alkoxy group of from one to six carbon atoms, a hydroxy group, hydrogen, an alkanoyloxy group, a benzoyloxy or substituted benzoyloxy group, a carbonate group or a carbamate group.

8. (Original) A dosage form of a pharmaceutical composition as claimed in claim 1 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine and lycoramine wherein the hydroxy group of galanthamine or lycoramine is replaced by an alkoxy group of from one to six carbon atoms, hydrogen, an alkanoyloxy group, a benzoyloxy or substituted benzoyloxy group, a carbonate group or a carbamate group.

9. (Original) A dosage form of a pharmaceutical composition as claimed in claim 1 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine and lycoramine wherein the N-methyl group of galanthamine or lycoramine is replaced by hydrogen, alkyl, benzyl or a cyclopropylmethyl group or a substituted or unsubstituted benzoyloxy group.

10. (Original) A dosage form as claimed in claim 7 wherein said acetylcholinesterase inhibitor is selected from the group consisting of galanthamine, lycoramine and analogs thereof wherein the methoxy group of such compounds is replaced by a hydrogen, hydroxy or alkoxy group of from two to six carbon atoms or an acyloxy group of from one to seven carbon atoms.

11. (Original) A dosage form as claimed in claim 7 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine or lycoramine wherein the methoxy group thereof is replaced by a mono or dialkyl carbamate or carbonate group wherein the alkyl groups contain from 1 to 8 carbon atoms

12. (Currently Amended) A dosage ~~form~~ form as claimed in claim 11 wherein the alkyl group or groups of said carbonate or carbamate groups comprise from 4 to 6 carbon atoms

13. (Original) A dosage form as claimed in claim 7 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine or lycoramine wherein the hydroxy group thereof is replaced by a mono or dialkyl carbamate or carbonate group wherein the alkyl groups contain

from 1 to 8 carbon atoms.

14. (Previously presented) A dosage form as claimed in claim 12 wherein the alkyl group or groups of said carbonate or carbamate groups comprise from 4 to 6 carbon atoms.

15.(Original) A dosage form as claimed in claim 7 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine or lycoramine wherein the methoxy group thereof is replaced by an aryl carbamate or carbonate group wherein said aryl group is selected from phenyl, naphthyl, substituted phenyl and substituted naphthyl groups wherein said substituent is selected from alkyl and alkoxy groups of from 1 to 6 carbon atoms, trifluoro methyl groups and halo groups.

16. (Original) A dosage form as claimed in claim 8 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine and lycoramine wherein the hydroxy group thereof is replaced by an aryl carbamate or carbonate group wherein said aryl group is selected from phenyl, naphthyl, substituted phenyl and substituted naphthyl groups wherein said substituent is selected from alkyl and alkoxy groups of from 1 to 6 carbon atoms, trifluoro methyl groups and halo groups.

17(Original) A dosage form as claimed in claim 8 wherein said acetylcholinesterase inhibitor is selected from the group consisting of galanthamine, lycoramine and analogs thereof wherein the hydroxy group of such compounds is replaced by a hydrogen or alkoxy group of from one to six carbon atoms or an acyl group of from one to seven carbon atoms.

18. (Original) A dosage form of a pharmaceutical composition as claimed in claim 7 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine and lycoramine wherein the hydroxy group of galanthamine or lycoramine is replaced by an alkoxy group of from one to six carbon atoms, hydrogen, an alkanoyloxy group, a benzoyloxy or substituted benzoyloxy group, a carbonate group or a carbamate group.

19.(Original) A dosage form as claimed in claim 7 herein said acetylcholinesterase inhibitor is galanthamine.

20.(Original) A dosage form as claimed in claim 1 wherein said acetylcholinesterase inhibitor is rivastigmine.

21.(Currently amended) A method of treatment of a patient suffering from a disease or condition in which it is desirable to administer a centrally acting acetylcholinesterase inhibitor which comprises administering a dosage form of pharmaceutical composition which comprises an effective amount of a centrally acting acetylcholinesterase inhibitor useful in the treatment of Alzheimer's disease selected from the group consisting of galanthamine, lycoramine, analogs of galanthamine and lycoramine rivastigmine said compounds having a half life of from one to eleven hours wherein the acetylcholinesterase inhibitor is formulated so as to delay its activity for a specified period of from four to twelve hours, wherein said administration is effected at a time prior to a patient's sleeping such that the activity of the cholinesterase inhibitor is delayed until after the patient has completed a period of sleep. .

22. (Original) A method of treatment as claimed in claim 21 wherein said patient is suffering from Alzheimer's disease.

23. Canceled

24. (Previously presented) A method of treatment as claimed in claim 21 wherein the composition is formulated to delay the activity of the acetyl cholinesterase inhibitor for a period of from six to nine hours.

25. (Previously presented) A method of treatment as claimed in claim 21 wherein

the composition is formulated to delay the activity of the acetyl cholinesterase inhibitor for a period of from eight to twelve hours.

26. (Original) A method of treatment as claimed in claim 21 wherein said acetylcholinesterase inhibitor has a duration of action of from 2 to 12 hours.

27. Canceled

28. (Original) A method of treatment as claimed in claim 21 wherein said acetylcholinesterase inhibitor is selected from the group consisting of galanthamine, lycoramine and analogs of said compounds wherein at least one of the methoxy, hydroxy or methyl groups of the galanthamine or lycoramine is replaced as follows:

the methoxy group by another alkoxy group of from one to six carbon atoms, a hydroxy group, hydrogen, an alkanoyloxy group, a benzoyloxy or substituted benzoyloxy group, a carbonate group or a carbamate group;

the hydroxy group by an alkoxy group of from one to six carbon atoms, hydrogen, an alkanoyloxy group, a benzoyloxy or substituted benzoyloxy group, a carbonate group or a carbamate group;

the N-methyl group by hydrogen, alkyl, benzyl, cyclopropylmethyl or a substituted or unsubstituted benzoyloxy group.

29. (Original) A method of treatment as claimed in claim 28 wherein said acetylcholinesterase inhibitor is selected from the group consisting of galanthamine, lycoramine and analogs thereof wherein the methoxy group of such compounds is replaced by a hydrogen, hydroxy or alkoxy group of from two to six carbon atoms or an acyloxy group of from one to seven carbon atoms.

30. (Original) A method of treatment as claimed in claim 28 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine or lycoramine wherein the methoxy group thereof is replaced by a

mono or dialkyl carbamate or carbonate group wherein the alkyl groups contain from 1 to 8 carbon atoms.

31. (Original) A method of treatment as claimed in claim 30 wherein the alkyl group or groups of said carbonate or carbamate groups comprise from 4 to 6 carbon atoms.

32. (Original) A method of treatment as claimed in claim 31 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine or lycoramine wherein the hydroxy group thereof is replaced by a mono or dialkyl carbamate or carbonate group wherein the alkyl groups contain from 1 to 8 carbon atoms.

33. (Original) A method of treatment as claimed in claim 32 wherein the alkyl group or groups of said carbonate or carbamate groups comprise from 4 to 6 carbon atoms.

34. (Original) A method of treatment as claimed in claim 28 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine or lycoramine wherein the methoxy group thereof is replaced by an aryl carbamate or carbonate group wherein said aryl group is selected from phenyl, naphthyl, substituted phenyl and substituted naphthyl groups wherein said substituent is selected from alkyl and alkoxy groups of from 1 to 6 carbon atoms, trifluoro methyl groups and halo groups.

35. (Original) A method of treatment as claimed in claim 28 wherein said acetylcholinesterase inhibitor is selected from the group consisting of analogs of galanthamine and lycoramine wherein the hydroxy group thereof is replaced by an aryl carbamate or carbonate group wherein said aryl group is selected from phenyl, naphthyl, substituted phenyl and substituted naphthyl groups wherein said substituent is selected from alkyl and alkoxy groups of from 1 to 6 carbon atoms,

trifluoro methyl groups and halo groups.

36. (Original) A method of treatment as claimed in claim 28 wherein said acetylcholinesterase inhibitor is selected from the group consisting of galanthamine, lycoramine and analogs thereof wherein the hydroxy group of such compounds is replaced by a hydrogen or alkoxy group of from one to six carbon atoms or an acyl group of from one to seven carbon atoms.

37. (Original) A method of treatment as claimed in claim 28 wherein said acetylcholinesterase inhibitor is galanthamine.

38. (Original) A method of treatment as claimed in claim 21 wherein said acetylcholinesterase inhibitor is rivastigmine.

39. (Original) A method of treatment as claimed in claim 21 wherein said acetylcholinesterase inhibitor is administered in conjunction with a compound that reduces its peripheral effects.

40. (Original) A method of treatment as claimed in claim 39 wherein said acetylcholinesterase inhibitor is administered in conjunction with a suitable dose of probanthine or robinul.

41. (Newly presented) A method of treatment of a patient suffering from a disease or condition in which it is desirable to administer a centrally acting acetylcholinesterase inhibitor which comprises administering a delayed release dosage form of a pharmaceutical composition which comprises an effective amount of a centrally acting acetylcholinesterase inhibitor selected from the group consisting of galanthamine, lycoramine, analogs of galanthamine and lycoramine rivastigmine said compounds wherein said dosage is formulated such that the half life of the activity of the acetyl cholinesterase inhibitor and the degree of delayed release are selected such that the formulation may be administered to the patient prior to the start of the patient's normal sleep time, acetyl cholinesterase inhibition

is delayed for the anticipated duration of said sleep time and the acetylcholinesterase inhibition resulting from such administration is reduced prior to the patient's next following normal sleep time and wherein administration of the drug is prior to an anticipated sleep time in order to achieve such results.